

Tidal chain reaction and the origin of replicating biopolymers¹

Richard Lathe

Pieta Research, PO Box 27069, Edinburgh EH10 5YW, UK
e-mail: rlathe@pieta-research.org

Abstract: Template-directed polymer assembly is a likely feature of prebiotic chemistry, but the product blocks further synthesis, preventing amplification and Darwinian selection. Nucleic acids are unusual because charge repulsion between opposing phosphates permits salt-dependent association and dissociation. It was postulated (Lathe, R. (2004). Fast tidal cycling and the origin of life. *Icarus* **168**, 18–22) that tides at ocean shores provide the driving force for amplification: evaporative concentration promoted association/assembly on drying, while charge repulsion on tidal dilution drove dissociation. This permits exponential amplification by a process termed here the tidal chain reaction (TCR). The process is not strictly contingent upon tidal ebb and flow: circadian dews and rainfalls can produce identical cycling. Ionic strength-dependent association and dissociation of nucleic acids and possible prebiotic precursors are reviewed. Polymer scavenging, chain assembly by the recruitment of pre-formed fragments, is proposed as the primary mechanism of reiterative chain assembly. Parameters determining prebiotic polymer structure and amplification by TCR are discussed, with the suggestion that Darwinian selection may have operated on families of related polymers rather than on individual molecules.

Received ■ 2005, accepted ■ 2005

Key words: DNA, evolution, Moon, nucleic acid, PCR, polymer, tidal chain reaction (TCR).

AQ1

Introduction

‘If it were given to me to look beyond the abyss of geologically recorded time ... I should expect to be a witness of the evolution of living protoplasm from not living matter ... I should expect to see it appear under forms of great simplicity, endowed ... with the power of determining the formation of new protoplasm from such matters as ammonium carbonates, oxalates and tartrates, alkaline and earthy phosphates, and water, without the aid of light’. In these words Thomas Huxley saw the emergence of life from simple prebiotic materials (Huxley 1870).

Haldane (1929) and Oparin (1953) proposed that the early ocean was a prebiotic soup, a rich diversity of organic and inorganic compounds for assembly into more complex precursors of true lifeforms. Miller and Urey addressed this by subjecting a mixture of ammonia, methane, hydrogen and water to heat and electric discharge, and saw the production of hydrogen cyanide, aldehydes and amino acids (see Miller (1953), Miller & Urey (1959) and Ferris (1987)). These can be further converted to nucleobases (from HCN and methane) and sugars (from HCHO) (Oro 1961; Sanchez *et al.* 1967; Sanchez & Orgel 1970; Fuller *et al.* 1972; Ferris & Hagan 1984; Shapiro 1995). In turn, polymerization can generate polypeptides and, notably, polynucleotides (Fox & Harada

1958; Schramm *et al.* 1962; Naylor & Gilham 1966; von Kiedrowski *et al.* 1989; Ferris *et al.* 1996; Luther *et al.* 1998).

The Watson–Crick alignment of precursors along a template strand, promoting chemical linkage to generate a complementary strand, is a compelling model for the origin of self-replicating systems, but a question arises regarding the driving force for association and dissociation. Template-directed polymerization is a dead-end. *‘Somewhere in this cycle work must be done, which means that free energy must be expended. If the parts assemble themselves on a template spontaneously, work has to be done to take the replica off; or, if the replica comes off the template of its own accord, work must be done to put the parts on in the first place’* (Blum 1957).

Thus, despite ample evidence suggestive of prebiotic polymerization, the evidence argues against cyclic copying in the absence of a further driving force. A previous proposal that rhythmic tidal flooding provided this driving force (Lathe 2004) is put into context. This paper attempts to provide a broad general overview; for in depth discussions of particular aspects the reader is addressed, where possible, to more detailed treatments.

Tidal chain reaction (TCR)

The association of two nucleic acid strands to form a double-helical structure, as in DNA, is determined not only by temperature, but also by ionic strength. This offers the

¹ Dedicated to Dr Harold F. Blum.

possibility that rhythmic changes in salt concentration brought about by dilution and drying permitted early biopolymers to associate (with template-directed polymerization) and dissociate (see below). This formally resembles the polymerase chain reaction (PCR), cyclic enzyme-mediated polymerization where repetitive changes in temperature (rather than ionic strength) drive exponential amplification. These two processes are contrasted below.

PCR

This technique is now widely used for DNA typing in forensic and genetic studies. PCR permits repetitive amplification of a DNA sample, in the presence of a thermostable polymerase enzyme and monomeric precursors, by oscillating the sample between two temperatures: at 50 °C single DNA strands direct complementary strand synthesis (doubling the molecule number), then on a shift to 100 °C the strands dissociate to permit the next round of synthesis. Multiple cycling produces exponential amplification, converting a single starting DNA molecule, *in vitro*, into 10^{15} identical molecules over 40 cycles (Mullis *et al.* 1986).

On the early Earth there is no evidence for substantial marine temperature cycling, although night-time cooling and day-time heating will have had some impact on surface (ocean) temperatures (discussed further below). Instead, the cycling of salt concentration may have provided a driving force (Lathe 2004).

TCR

Modern nucleic acids are unusual because, unlike most biopolymers, the association of the two complementary strands is promoted by increasing ionic strength (Schildkraut & Lifson 1965). This property reflects the structure of the double helix, where strong charge repulsion between phosphate groups is neutralized by increasing salt concentration (Fig. 1).

As a consequence, nucleic acids may have replicated through tide-driven drying and dilution. Elevated salt concentrations produced on drying promote association of single strands to form duplex structures; then at low ionic strength (e.g. following tidal inflow or rainfall) the duplex is driven to dissociate.

By the same mechanism, salinity increase is likely to favour association of monomeric precursors with a preformed template strand: tight alignment favours the formation of covalent links between adjacent monomers (Fig. 2). Partially anhydric conditions also encourage non-enzymatic polymerization through dehydration reactions (Schramm *et al.* 1962; Calvin 1969; Weber *et al.* 1977).

Two rules for the origin of life

For primitive biopolymer replication to emerge from tidal cycling, two rules are likely to apply. First, because precursor concentrations were certainly limiting, polymerization must take place during the drying phase when precursors are most abundant; partial anhydric conditions also favour chemical bond formation. Second, as noted by Blum (1957), if assembly takes place on drying, work must be done to remove




Fig. 1. Salt-dependent association and dissociation of nucleic acids.




Fig. 2. Salt increase on drying promotes both (a) duplex formation and (b) monomer alignment for polymerization; salt removal on dilution then drives dissociation.

the copy so as to permit amplification: dissociation must therefore take place on dilution. It is suggested here that only molecules similar to modern nucleic acids fulfil these two criteria.

Reconsideration of parameters

Emphasis is placed on the rapidity of tidal cycling and the ionic conditions of the early ocean. These and other relevant parameters are revisited.

The importance of rapid cycling: inherent instability of polymers

The speed of cycling is important because of product instability. Nucleobases are generally stable at high temperatures (half-lives in the order of weeks or more at 100 °C for adenine and cytosine), ribose is reported to have a half-life of only 73 minutes at the same temperature (Larralde *et al.* 1995; Miller & Lazcano 1995; Levy & Miller 1998), although deoxyribose (lacking the 2' hydroxyl of ribose) is more stable.

However, the instability of a polymer increases linearly as a function of length. For example, if under Archaeal conditions the chemical half-life (leading to chain scission) of a monomer in primitive nucleic acid is 1 week, the half-life of a 100-mer strand is only ~ 3 h. Ultraviolet irradiation and radioactivity compound the problem.

Because absolute biopolymer number depends on the rate of assembly by *de novo* polymerization (or by scavenging, see below) *versus* the rate of fragmentation/ degradation by thermochemical and other processes, rapid cycling may be critical to permit biopolymer accumulation.

A second perspective concerns the total tidal volume subjected to cyclic drying and dilution. The absolute number of polymers may be critical if a further rare event or events are required to generate bona-fide lifeforms. Candidate rare events include the origin of coding and micellar inclusion to generate cellular life. If these events are rare, the likelihood of occurrence will depend on the absolute number of opportunities for their occurrence. The emergence of true lifeforms is favoured by, but not contingent upon, large and rapid tidal cycling.

There is a downside to rapid cycling: because evaporation is time-dependent, rapid cycling will also limit the extent of water loss from large volumes. Simulations will be required to address the constraints that cycling speed places on evaporative concentration.

Early oceans

Oceans first formed when surface temperatures fell below 100 °C. Although there was a period of intense bombardment (Hartmann 1980), leading to an unstable surface and potentially extinguishing prebiotic evolution several times (Maher & Stevenson 1988), by -4 Ga it is thought that stable water cover existed at the terrestrial surface. Ocean temperatures in the range 50–100 °C are plausible, stabilized by high atmospheric CO₂ content (Kasting & Ackerman 1986).

The early salinity of the ocean remains a matter of debate. Concentrations have remained fairly constant over the last 1 Ga or more (Knauth 1998), but before that they may have been higher. The conventional view is that volatiles outgassed rapidly and, because sequestration could not take place until continent formation (-2.5 Ga), the early ocean must have been highly saline (Knauth, Personal communication). A recent review of this topic concluded that early oceans were hot and super-saline, with a NaCl concentration somewhat in excess of present-day levels (Knauth 2005).

AQ2 Early salinity is an important issue because excess [NaCl] can block TCR amplification (see below), challenging an oceanic origin for replicating biopolymers. Nevertheless, empirical data for (or against) a highly saline ocean are limited. An alternative scenario is that the high chemical reactivity of outgassed chlorines (and other elements) would have led to rapid deposition of solid phase chlorides prior to ocean formation. These salts would only have commenced dissolution on condensation of the oceans. Under this interpretation the salt concentration could have been lower than at present. To speculate: if it took 1 Ga for ocean salt to

rise to near present day values (460 mM NaCl) and only 0.1 Ga for life to evolve, the [NaCl] range in the first 100 Ma might have been between 0 and 46 mM (values consistent with the association/dissociation of long polymers, see below). However, given conflicting estimates and the crucial importance for biopolymer replication, the issue of early salinity needs to be resolved with urgency.

Six ions comprise $\sim 99\%$ of salts in the present-day ocean: chloride (Cl⁻), sodium (Na⁺), sulfate (SO₄²⁻), magnesium (Mg²⁺), calcium (Ca²⁺) and potassium (K⁺), with NaCl by far representing the major component. Carbonates and ammonium ions may have been far more abundant, reflecting CO₂ and NH₃ in the Hadean/Archaean atmosphere, but phosphorus ions (principally phosphates), key ingredients of modern nucleic acids and their precursors, are relatively insoluble (present-day concentrations up to 1 mM), although pyrophosphates appear to be more soluble.

Early tides

The frequency of early tidal ebb and flow could have been far faster than at present, but the evidence is contradictory. Fast rotation of the early Earth has been associated with the impact event that produced the Moon, and since that time the Moon is presumed to have receded from the Earth, with concomitant slowing of the rotation rates of both bodies. Darwin (1879) suggested an early terrestrial rotation period of under 6 h (discussed in Brush (1986); see also Hartmann (1999)).

Relevant empirical data have been derived from the accurate measurement of the fast recession rate (3.82 cm yr⁻¹) of the Moon over recent decades (Dickey *et al.* 1994) and analysis of striations in paleobiological residua (fossil corals, bivalves, stromatolites and tidalites) from the last 1–2 Ga (Panella 1975; Scrutton 1978; Lambeck 1978, 1980; Vanyo & Awramik 1982; Sonett *et al.* 1996; Williams 1997, 2000; Kvale *et al.* 1999). Regression curves from these empirical data (dependent both on the length of day and lunar orbital rate) suggest steep and steady recession and orbital decay in the last 1.5 Ga, pointing to cataclysmic proximity between -1.5 and -3 Ga, an event for which there is no evidence (Walker & Zahnle 1986). It is possible that the Earth–Moon pair has decayed in two phases, first slowly to about -2 Ga, followed by more rapid decline to present-day values (Krasinsky 2002). With limited and contradictory data it is unsafe to attempt to retrodict relative rotation at -3.9 Ga with any accuracy. Despite this, the relative Earth versus Moon rotation frequency was probably faster than at present, although whether 6 or 16 h (with tides every 3–8 h) is not yet known (see Varga *et al.* (2005) and Lathe (2005) for further debate).

Early shores and tidal flooding

At present, 70% of the terrestrial surface is covered by ocean; in the Archaean the Earth is felt to have been far flatter and ocean coverage may have exceeded 90%. The extent of early tidal flooding is therefore subject to great uncertainty. Factors determining tidal height and extent are the hindrance

(i.e. size) afforded by land mass and coastal geometry, with harmonic resonances playing a major role (Butikov 2002). Present day estuarine tides of 16 m are recorded in shallow water at the Bay of Fundy (Nova Scotia), where tidal focusing takes place, but in Iceland (a largely volcanic outcrop in the open ocean, perhaps representative the Archaean land surface) the tides are much reduced in magnitude (typically less than 1 m), pointing to smaller tides in that epoch.

However, it may not be assumed that the total liquid water volume has remained constant. A greater proportion may have been present as vapour in a warmer epoch, exposing more land. There are further complexities regarding the chemistry and rate of water exchange with minerals. The early Moon was probably much closer to the Earth, with far greater tidal forces, but the extent of tidal flooding (even at a defined Earth–Moon distance) is non-obvious, and numerical tidal modelling will be required, not so far performed.

DNA dissociation at low ionic strength

A core contention of the tidal cycling theory is that the two strands of DNA associate at high ionic strength, but separate at low ionic strength, largely due to charge repulsion between opposing phosphate groups. This association and dissociation is inherent to the Schildkraut–Lifson (1965) relation and extended in other analyses (Thomas & Dancis 1973; Wetmur & Davidson 1986; Wetmur 1991):

$$T_m = 81.5 + 16.6 \log M + (0.41\%G + C) - 820/l$$

where T_m (melting temperature) is the 50% dissociation temperature, M is the monovalent cation (Na^+) concentration, %G + C is the percentage composition of the duplex in guanosine and cytosine and l is the length of the duplex (see Lathe (1985)).

However, it is accepted in the molecular biology laboratory that even small duplexes (e.g. 20-mer complementary oligonucleotides) can remain associated at moderate temperatures and low salt concentrations. This raises the question of whether DNA truly dissociates, at room temperature, on reduction of [NaCl].

The stability of a nucleic acid duplex is commonly expressed in terms of melting temperature or T_m , the temperature (in °C) at which 50% of the sample is dissociated. This has two non-obvious consequences.

First, T_m becomes dependent on the length of the duplex. This is because association/dissociation at temperatures close to the T_m is dynamic: subdomains dissociate while other regions remain associated ('breathing'). Because the complementary strands are physically retained in proximity, the chains do not diffuse apart, favouring reassociation. A negative reciprocal function (e.g. $820/l$ (Lathe 1985) or $660/l$ (Hames & Higgins 1985), where l is the length of the duplex) is appended to the Schildkraut–Lifson relation to take account of this.

Second, a rise in strand concentration strongly increases the apparent T_m , as expected for a conventional two-order reaction. High concentrations of oligonucleotide primers and




Fig. 3. Dissociation (melting) temperature, T_m , of a 100-mer DNA molecule (50% G + C content) as a function of ionic strength; calculated from the Schildkraut–Lifson relation as modified (Lathe 1985).

linkers are often employed in the laboratory, raising T_m by as much as 50 °C above the value predicted from the Schildkraut–Lifson relation.

Melting temperatures at low strand concentrations, such as those expected in the prebiotic ocean, are much reduced. A standard DNA 60-mer oligonucleotide duplex dissociates, in 10 mM NaCl, at 55–70 °C (Schildkraut & Lifson 1965) or below, depending on base composition (the standard Schildkraut–Lifson equation omits the concentration dependency). Dissociation of a perfectly matching 169-nucleotide DNA duplex in 10 mM NaCl at 61.5 °C has been empirically demonstrated (Rose *et al.* 2002). At further reduced salt concentrations it was observed (in a study on DNA stretching) that if NaCl was eliminated from the buffer, typically parts of or even the entire molecule dissociate before overstretching is applied (Clausen-Schaumann *et al.* 2000). One must conclude that, at 50 °C and low [NaCl], protonucleic acids will have dissociated into separate chains, only reassociating on evaporative concentration.

Divalent cations

Salt [NaCl] is not the only variable: trace divalent ions including Mg^{2+} can stabilize nucleic acid duplexes to a far greater extent than the same concentration of Na^+ (Williams *et al.* 1989). In an extended study, Nakano *et al.* (1999) reported that Mg^{2+} , Ca^{2+} , Mn^{2+} and Ba^{2+} are far more effective stabilizers than Li^+ , Cs^+ , Na^+ or K^+ . The melting temperature of a short DNA duplex was disproportionately increased by 4 °C when 10 mM Mg^{2+} was added to a solution of 100 mM NaCl (Nakano *et al.* 1999). However, Mg^{2+} is not abundant in seawater (presently ~40 mM) and early concentrations may have been very much lower. The trace concentrations inferred for the Archaean ocean (up to 2 mM, paralleling [NaCl]) are probably insufficient to exert a major effect (although, as with NaCl, significant concentrations are expected on drying).

How do ions stabilize nucleic acids?

Monovalent cations bind to and neutralize charges on opposing phosphate groups. However, the interactions of ions (including water) with DNA and RNA are complex. Phosphates, sugars and the sides of the base pairs are exposed to and interact with solvent in both the minor and major grooves (Westhof 1988; Auffinger *et al.* 2004; Schneider *et al.* 1998). Interactions are not restricted to cations, and include anionic chloride and sulfate (Auffinger *et al.* 2004). Ion exposure modulates both charge and base stacking interactions – in the Watson–Crick duplex, a major stabilizing force is the juxtaposition of parallel nucleobases (or base pairs), attributed to orbital overlap. Interacting ions modulate this interaction: stabilization and destabilization of the double helix is more complex than simple charge neutralization.

In partly anhydric conditions, brought about either by drying or by organic solvent replacement, the standard B-form duplex undergoes a series of transitions to more compact forms including the A-form helix (Franklin & Gosling 1953; Piskur & Rupprecht 1995) but which conserve base-pairing specificity. If evaporation routinely proceeded to near-dryness and precursor assembly took place under these conditions, the A-form could be as relevant as the B-form.

Chaotropism

As we have seen, increasing [NaCl] increases DNA stability according to a log function over the range 0–1 M (Schildkraut & Lifson 1965). At even higher concentrations (>2 M) the converse effect is seen, with a reduction in stability. This is ascribed to the chaotropic effects of high ion concentrations. Chaotropicity is a function of ion size (and concentration), where small ions (Na, Cl, Li) exert minimal effects while large ions (trichloroacetate, guanidinium, perchlorate) dissociate rather than stabilize nucleic acid duplexes (Hamaguchi & Geiduschek 1962; Haung 1968; Ott *et al.* 1975). Less ionic molecules (urea, formamide, imidazole) can also dissociate nucleic acids (Cox & Kanagalingam 1968; McConaughy *et al.* 1969; Kourilsky *et al.* 1970, 1971; Casey & Davidson 1976; Eli *et al.* 1999), but only at high concentrations (e.g. 1–5 M or 50% w/v).

The abundance of chaotropic molecules in the prebiotic ocean has not been assessed. Urea is formed in significant quantities in prebiotic simulations (Schlesinger & Miller 1983) and, if tidal recession permitted evaporation to near-dryness, it is plausible that association of early nucleic acids was followed by a dissociation phase at extremely high solute concentrations. The possibility that such pre-dissociation might influence strand separation on subsequent tidal dilution deserves investigation.

Condensation versus chaotropicity

In solution, DNA will spontaneously collapse and aggregate by a process termed condensation. This usually requires cations with a charge +3 or greater (Bloomfield 1991) and ions used in condensation studies include the naturally-occurring




Fig. 4. Formation of a triple-helical structure by the addition of a third strand to the standard double helix (schematic).

polyamines spermine and spermidine (Gosule & Schellman 1976; Burak *et al.* 2003). Large (potentially chaotropic) ions of positive/mixed charge also cause collapse by simultaneously neutralizing surface phosphates while, at the same time, permitting interactions between adjacent helices. Collapse is inhibited by potent chaotropic ions such as trichloroacetate (Ott *et al.* 1975).

Whereas divalent metal ions do not normally provoke condensation, they can do so in water–ethanol mixed solution (Arscott *et al.* 1990b). High ethanol concentrations are unlikely in the prebiotic ocean, but it is plausible to suggest that removal of water by evaporation may have left primitive biopolymers in a milieu enriched in non-volatiles including oils and salts, conducive to condensation, leading to large particulate aggregates. One feature of such a collapse is the formation of toroidal structures (Eickbush & Moudrianakis 1978; Arscott *et al.* 1990a). These could presage the emergence of circular chromosomes (see also below) and further study is warranted.

Did triple helices play a role?

One peculiar feature of modern day DNA is the ability to form triple- (rather than merely double-) helical structures. This is because the standard double helix contains a lateral gap, the ‘major groove’, into which a further strand can fit (Fig. 4). The sides of the nucleobases (base-pairs) are exposed in the groove, allowing them to form unconventional (non-Watson–Crick) pairs with a third strand.

Triple helices are stabilized by many factors that stabilize the duplex, including polyamines (Musso & Van Dyke 1995; Musso *et al.* 1997; Saminathan *et al.* 1999) and divalent cations (Blume *et al.* 1999; Floris *et al.* 1999). However, monovalent cations inhibit triple-helix formation (Cheng & Van Dyke 1993; Singleton & Dervan 1993; Plum *et al.* 1995; Floris *et al.* 1999), arguing against the possibility that triple-helical structures were a major feature of tide-driven prebiotic replication. This conclusion would need revision if it can be shown that polyamines or equivalent molecules were abundant in the early ocean, a possibility that is not unlikely.




Fig. 5. Polymer scavenging. (a) Association of monomers with a template strand leading to chemical polymerization and dissociation on dilution. (b) The same process whereby short pre-formed polymer fragments align on a template strand for chemical ligation. (c) If longer polymers align in this way, an overall increase in chain length is expected.

Getting started: minerals

TCR is proposed as a scheme whereby primitive polymers directed their own replication through the ordered assembly of complementary monomers into a complementary strand.

The first polymers may have assembled by chance, but assembly can be catalysed by mineral surfaces (Bernal 1949, 1967). Clays and silicates promote alignment and polymerization of precursors; the structure of the earliest nucleic acids may have borne an imprint of the facilitating mineral surface (Cairns-Smith 1982, 2001). Surface association facilitates polymerization in several models (Ferris *et al.* 1983, 1996; Luther *et al.* 1998; Ertem & Ferris 1996, 2000; Orgel 1998; Liu & Orgel 1998). TCR-cycling may extend to mineral catalysis, as nucleic acids will adsorb to clays/silicates at high salt concentration, facilitating polymer synthesis, but not at low salt concentration, permitting dissociation on dilution (Vogelstein & Gillespie 1979; Mizutani & Narihara 1982).

Copy number and length increase

Tidal cycling predicts an exponential increase in the number of prebiotic polymers structurally similar to nucleic acids. It also predicts an increase in chain length. Just as two monomers are inferred to align on a template strand during the drying phase, prompting chemical bond formation between them (Fig. 5(a)), two part-formed polymers are likely to juxtapose by base-pairing with a template strand (Figs 5(b) and (c)), also favouring covalent linkage.

Ligation-mediated PCR

This polymer ligation process is formally similar to the PCR but, instead of using an enzyme to direct polymerization,

linkage between two abutting oligonucleotides is driven through the addition of the ligase enzyme (Wu & Wallace 1989; Barringer *et al.* 1990). Combined ligation and polymerase-driven reactions have been performed (Bi & Stambrook 1997). The efficiency of this reaction supports scavenging (below) as a realistic process.

Ligation TCR: polymer scavenging

Ligation of short pre-formed polymers, a process termed 'polymer scavenging' here, is an alternative mechanism of complementary strand generation in a prebiotic context (Fig. 5(b)). Non-enzymatic ligation and amplification of oligonucleotides with suitably activated end-groups is efficient (Xu & Kool 1999; Gao & Orgel 2000). Given that the rate and affinity of association of short oligonucleotides with a complementary strand is far higher than for monomers with the same template, the process of complementary strand formation by scavenging (rather than *de novo* polymerization) is likely to have provided the primary mechanism for reiterative chain assembly. As in Figs 5(b) and (c), scavenging leads to an increase in both copy number and mean chain length. This process also opens the door to Darwinian selection in the sense where 'successful' polymers can exploit fragments produced by degradation of 'unsuccessful' molecules.

Circularity

A consequence of polymer scavenging is the generation of circular forms. When overall polymer abundance is low, the effective concentration of one end of a molecule versus the other exceeds the concentration (expressed in ends ml^{-1}) of other molecules and circularization becomes more likely than

extension (Jacobson & Stockmayer 1950; Dugaiczuk *et al.* 1975; Legerski & Robberson 1985). This effect may be enhanced by agents that promote the formation of condensed toroidal structures (see above). The formation of circular molecules is disadvantageous to prebiotic replication because the two strands are intertwined and dissociation on dilution is prevented. However, subsequent thermochemical breakage releases strand fragments for scavenging.

Caveats: primitive polymers

Were precursors sufficiently abundant to permit replication of nucleic acids? Although some studies indicated that precursors including phosphate-activated nucleosides (Gulick 1955; Schramm *et al.* 1962) were present, concentrations may have been low (Bernal 1960; Hull 1960). The relative instabilities of cytosine (Levy & Miller 1998) and ribose (with critically unstable 2'-3' adjacent hydroxyls) are cases in point. Prebiotic conditions were not favourable for the accumulation of these key ingredients (Shapiro 1988, 1995, 1999).

There are two routes out of this impasse. First, prebiotic conditions may not have been quite so hostile for the production of precursors.

Radioactivity is a significant source of organic molecule formation. Irradiation of methane, CO and CO₂ can efficiently generate alcohols and long-chain polymers, while radioactive minerals are particularly susceptible to hydrodynamic sorting and concentration on the margins of oceans (this has been reviewed by Parnell (2004)). These processes would favour the accumulation of complex organics in tidal regions.

Cyclic drying also provides a rather more favourable chemistry for precursor assembly. Anhydric conditions promote assembly by chemical dehydration reactions (Schramm *et al.* 1962; Calvin 1969; Weber *et al.* 1977); although challenged (Shapiro 2002), cytosine may be efficiently produced at 100 °C under dry-down conditions (Nelson *et al.* 2001).

Second, simpler and more abundant components may have preceded both DNA and RNA (Joyce *et al.* 1987; Shapiro 1995).

Base structure

Adenine is commonly generated in prebiotic simulations (Oro 1961; Sanchez *et al.* 1967); both adenine and guanine are stable at 100 °C (Levy & Miller 1998) while pyrimidines tend to be formed in low amounts and are less stable. Such considerations suggest that an all-purine nucleic acid may have predated the purine-pyrimidine pairing of both DNA and RNA (Wachtershauser 1988); a purine-only nucleic acid does not appear to be inconsistent with the emergence of coding.

Sugars

Low yields of ribose and deoxyribose in prebiotic experiments, coupled with the inherent instability of ribose, suggests the possibility of alternative sugars or related molecules as components of protonucleic acids, including glycerol, erythritol and acrolein (Joyce *et al.* 1987). Acrolein synthesis

occurs under simulated prebiotic conditions and chemical reactivity with nucleobases has been demonstrated (Cleaves 2002). Polymers based on the sugar threose have been shown to pair with both RNA and DNA (Schoning *et al.* 2000).

Phosphates and activating groups

Internucleoside phosphate groups provide the strong charge repulsion needed for TCR, but also reflect precursor activation (triphosphate in the case of present-day DNA and RNA). The incubation of modern nucleoside (DNA precursor) triphosphates at 74 °C with thermostable enzyme, in the absence of any template, has been shown to give rise to long double-stranded polymers (Ogata & Miura 1997); one must note that enzymes conventionally only catalyse reactions that would otherwise take place (although more slowly) in their absence. Therefore, one may presume that nucleoside triphosphates spontaneously convert to polymeric forms.

Nevertheless, there is a debate regarding the likely availability of phosphate- or polyphosphate-activated nucleosides. Solubility problems have been noted (Gulick 1955) and polyphosphate concentrations may fall short of those required for precursor activation (Keefe & Miller 1995). Nevertheless, volcanic activity is a rich source of P₄O₁₀, converting to polyphosphate on solution in water (Yamagata *et al.* 1991). P₂O₅ heated in anhydric organic solvent generates a polyphosphate ester with wide reactivity for organic molecules including nucleosides (Schramm *et al.* 1962). Phosphates in the presence of urea afford a further route to nucleoside activation (Lohrmann & Orgel 1971).

While the extent of phosphate activation is not yet known, *in vitro* studies have employed nucleosides 5'-activated with phosphorimidazolide (Imp): these can polymerize efficiently *in vitro*, a reaction stimulated by divalent cations and mineral surfaces (Lohrmann *et al.* 1980; Ding *et al.* 1996; Ertem & Ferris 1996; Ertem 2004). However, the existence of prebiotic Imp-activated nucleosides remains questionable.

If significant polyphosphate concentrations are questionable, one might speculate that other chemistries could have been employed. Unlike phosphate, sulfate is a major ion in the present-day ocean. Early biopolymers could have exploited a diversity of mixed phosphorus/sulphur/oxygen activating groups (and linkages). Phosphorothioates and phosphorodithioates retain the hybridization properties of standard DNA (Jaroszewski *et al.* 1996), while phosphorothioate linkages retain the charge repulsion property of the phosphodiester linkages in DNA and RNA. Other possible derivatives include sulfonyl (replacing phosphorus by sulphur) and boranophosphate linkages. Mixed sulphur-phosphorus chemistry in PAPS (3'-phosphoadenosine 5'-phosphosulfate) is exploited in present-day organisms. Nucleoside 5'-thio-sulfates and 5'-thiophosphates have been described (Golos *et al.* 2001), but without any evidence that they can participate in polymerization reactions. This area warrants further thought.

Nevertheless, phosphate activation remains the strongest contender. Schramm *et al.* (1962) reported remarkable synthesis of a phosphate-activated nucleoside by drying sugars



Fig. 6. Electron micrograph ($\times 60\,000$) of synthetic polyadenylic acid: from Fig. 1 of Schramm *et al.* (1962).

and adenine with polyphosphate ester. Further incubation of nucleotides with polyphosphate ester and pyridine at $50\text{--}60\text{ }^{\circ}\text{C}$ led to polymeric forms (Fig. 6) that were sensitive to nucleases.

Unconventional backbone structures

Polyamide or peptide nucleic acids (PNAs) fully eliminate the sugar phosphate moieties, replacing them by pseudo-peptide (polyamide/N-(2-aminoethyl)-glycine) groupings. PNAs have great affinity for natural nucleic acids, contributing to stable double-stranded molecules (Nielsen *et al.* 1991; Egholm *et al.* 1993), and PNA has been proposed as an alternative structure for the first replicating biopolymers (Nelson *et al.* 2000). Although the complementary-strand coding properties of such molecules are intact (Bohler *et al.* 1995), the lack of a large negative charge replacing the phosphate group prevents dissociation at low ionic strength. PNA-like molecules would seem unable to replicate by tidal cycling.

Chirality and catalysis

A major obstacle for all of the theories addressing the origin of replicating biopolymers concerns chirality. Chemical generation of precursors systematically results in a mixture of enantiomeric forms (D versus L), but only one is employed in present-day polymers (L amino acids in polypeptides, D sugars in nucleic acids; see Sandars (2005) for a review).

This problem is so far unresolved. However, it is well known that handedness is copied in crystalline forms: if mineral surfaces directed early polymerization it is likely that polymers were either D or L and rarely a mixture (see Cairns-Smith (1982)). For peptides, it has been argued that mixed L- and D-containing polymers will fail to form stable tertiary structures, favouring the accumulation of homochiral sequences (Brack 1987). By the same token, mixed (D and L) oligonucleotides may fail to adopt a stable double-helical structure. If this is so, strand-directed polymerization and scavenging could generate co-existing but separate homochiral D- and L-sugar forms.

Second, like RNA (Altman 1990), early nucleic acids (as single strands) may have folded to produce enzymatically-active molecules. This is enhanced under elevated temperature and cation concentrations (e.g. Peracchi (1999) and Takagi & Taira (2002)) typical of TCR. A catalytic molecule capable of discriminating between D and L sugars at

synthesis, assembly or isomerization could provide a plausible route to ‘only-D’ polymers.

Sequence motifs persist independently of exact molecular structure

The information content of a prebiotic polymer is largely only dependent on the base-pairing identity/specificity and position of the nucleobase. Therefore, a sequence motif may be conserved through successive replications even if the exact backbones and base structures are not.

We should consider the possibility that evolution operated on families or populations rather than on individuals. DNA can show catalytic (enzymatic) activity (Breaker 1997), but the more flexible RNA chain appears better adapted. If a parent molecule (e.g. DNA) is stable but catalytically inert, while daughters (e.g. RNA) are unstable but effective catalysts, selective pressures might operate on a mixed family of related polymers rather than on individual molecules. Coexistence might contribute to resolution of the DNA versus RNA debate (Dworkin *et al.* 2003).

Cyclicities: further reflections

Cyclic alteration to the local physicochemical environment affords a solution to Blum’s (1957) conundrum. Tides, proposed here as a major player, are one of several cyclicities, each with potential to drive association and dissociation.

Macro- versus micro-scale

Tidal flooding addresses pools on ocean margins, with large volumes of incoming water subsequently evaporating. This implicit assumption may be too restrictive, as parallel processes can easily occur on a micro-scale. The flooding of beaches leads to hydration of particulate sands; these then dry progressively on tidal ebb. Here the volumes of water surrounding each particle are much smaller, but the same processes will operate, although perhaps faster at the micro-scale. The total cycling volume is comparable to the tidal pool scenario.

Wave action

Ocean waves, the result of wind action on surface waters, could provide a mechanism for cyclic flooding. However, the cycling speed (seconds to minutes) is likely to be far too fast to allow significant physicochemical changes, for instance drying or heating/cooling. Volcanic and impact-driven (Tsunami) flooding merits consideration.

Spring and neap tides

Tidal ebb and flow is a major source of cycling, but tides are not uniform and follow higher-order cyclicities. The alignment of the Sun and Moon generates spring tides (unrelated to season) that are higher than the mean high tide; a week later (when the Sun and Moon are at 90°) their tidal influences partially cancel out, producing neap (lowest) tides (*neap*, Saxon, lacking or scanty; *nipflod*, Danish). Although of lower

periodicity, spring tides provide for more extensive flooding and a protracted drying phase.

Heat cycling: night/day

Terrestrial rotation provides a mechanism for cyclic heating and cooling; this was suggested as a possible driving force for the emergence of replicative molecules (Blum 1957). However, circadian changes in temperature of the present-day ocean are restricted to shallow areas; the same is likely to be true of the Hadean/Archaean ocean. Moreover, such cycling is not conducive to the inferred replication of prebiotic polymers, as heating (that dissociates duplex structures) counters association driven by an increase in salinity. A stagnant pool (without significant evaporation) cycling between 50 °C (night time) and 100 °C (day time) could afford PCR-type association and dissociation, but the absence of precursor concentration and regeneration on drying affords an unattractive scenario.

Rains and dews: TCR

Present-day clouds and periodic intense precipitation are contingent upon particulate matter suspended in the atmosphere. It is possible that the early atmosphere was, compared with the present day, relatively free of such particles, although volcanic activity would clearly contribute. If sporadic rainfall is questionable (and it may not be), a more likely scenario could involve condensation of water (dew) during the night-time phase with day-time re-evaporation. This process, if extensive, produces TCR cycling – condensation results in dilution (promoting the dissociation of prebiotic duplexes, and subsequent evaporation produces an increase in salt and precursor concentration (fostering the assembly of prebiotic duplexes).

Multivalent cycling

When the tidal cycle is out of phase with circadian cycling, the combination produces a faster cyclicality than either alone and deserves study. Tidal and circadian rhythmicities could combine to provide multivalent cycling of both temperature and salinity (simultaneous PCR and TCR). The cycle hot/drying, warm/dilute, warm/drying, hot/dilute would permit amplification of polymers with a structure similar to nucleic acids.

Submarine vents

Deep-sea hydrothermal vents (Corliss *et al.* 1979) have been suggested as a possible site for the origin of life. In support, extreme thermophiles are among the earliest cellular organisms (Woese *et al.* 1990; Woese 2000) and such hot vents provide a honey-pot of chemical energy and complexity (Ferris 1992; Shock 1996; Nisbet & Sleep 2001; Reysenbach & Shock 2002). Vent-driven convective cycles of heating and cooling could simulate PCR conditions and oligomerization of amino acids has been demonstrated in a simulation (Imai *et al.* 1999). Nevertheless, the hydrothermal hypothesis has been questioned because the extremely high temperatures decompose organic molecules (Miller & Bada 1988); less extreme temperatures at adjacent sites may have been more

conducive (MacLeod *et al.* 1994). A greater problem concerns the dilution factor. Heat dissociation of prebiotic polymers in the locality of the vent is followed by dilution (as hot liquid rises and cools), opposing reassociation and assembly. It is not felt that open hydrothermal vents and macro-scale convection afford a promising candidate for chain amplification reactions driven by either salinity or heat cycling, but other mechanisms may operate.

PCR by laminar convection

PCR is performed in the laboratory by imposing rapid temperature shifts; this is argued here to be an unlikely scenario at the origin of life. However, two groups have demonstrated that thermal (PCR) cycling can take place in a steady temperature gradient (Krishnan *et al.* 2002; Braun *et al.* 2003). In a typical experiment, a fluid-filled 5 × 1 mm disc was heated in the centre using an infrared lamp, inducing stable laminar (toroidal) convection. DNA dissociates in the hot centre, from where it is transported to the periphery for copying mediated by thermostable polymerase included in the reaction mixture. Further convection transports the DNA back to the centre. The amplification achieved was comparable to conventional heat-cycling PCR (Braun *et al.* 2003). Krishnan *et al.* (2002) used a vertical sandwich of similar size heated from below and cooled at the top to achieve a comparable result.

Such closed-chamber PCR devices profit from a phenomenon known as thermophoresis (or thermodiffusion) where molecules are repelled from a heat source and move down a hot–cold gradient. A 50 °C temperature gradient led to a 1000-fold enrichment of a 5 kb DNA in the cold zone (Braun & Libchaber 2002).

Braun & Libchaber (2004) suggested that such mechanisms operated at the origin of life, notably in association with hydrothermal vents. They postulated that porous rocks could have provided the small (<1 cm) chambers required, with temperature gradients generated by the contrast between the hydrothermal vent outflow (>300 °C) and ocean bottom seawater (perhaps as low as 4 °C).

Disadvantages with these scenarios at the origin of life include the need for a closed chamber and a steep temperature gradient, and the absence of any mechanism for concentrating and generating precursors. Thermophoresis could have contributed to this mechanism, but smaller molecules (27 nucleotides) were less than two-fold concentrated under the conditions described by Braun & Libchaber (2002). Nevertheless, thermal cycling remains a viable contender.

Salinity: tides versus dews

The salinity of the early ocean is not known, but some estimates have suggested that it could have been higher than present-day values (in excess of 0.5 M NaCl) (Knauth 2005). This affords a major barrier for chain amplification by dilution and concentration – even short duplexes (e.g. 100-mer) only dissociate at elevated temperatures at this NaCl concentration (>80 °C; see Fig. 3) and dissociation by tidal

inflow becomes less plausible. It remains possible that early [NaCl] was far lower, but if high early salinity should be confirmed this would argue against tidal inflow as the drive for dissociation. However, TCR amplification (defined as cyclic dilution and evaporation) could exploit condensation (dews and rains), relocating the origin of replicating biopolymers away from coastal areas to inland lakes and pools.

Freshwater tides

Even at high ocean salinity, tidal salt cycling takes place in estuaries. Outflowing freshwater is backed up by tidal inflow, only to be released (with drying sands and pools) on tidal ebb. Flood–ebb cycles can affect river levels for hundreds of kilometres inland (with far inland tidal ranges up to 4 metres being recorded in St. Lawrence). Within the upper reaches of the Bay of Fundy salinities vary from nearly marine to fresh within a single tidal cycle (Kvale, Personal communication). Estuarine tides afford an alternative location for prebiotic TCR cycling.

Discussion

The amplification of a primitive biopolymer requires repetitive assembly and dissociation, a process blocked under steady-state conditions (Blum 1957). Tidal cycling allows this process to take place. The importance of coastal areas and evaporative drying has been recognized previously (Blum 1957, 1962; Bernal 1961, 1967). Cyclic drying and dilution (TCR) affords a mechanism for the repetitive assembly and dissociation of primitive biopolymers resembling nucleic acids (Lathe 2004). High early ocean salinity (if confirmed) would inhibit this process, but would point instead to inland lakes and pools (via circadian condensation and precipitation) and estuaries.

Simple considerations suggest two rules. Association and polymerization must take place during the drying/concentration phase, when precursors are most concentrated; the duplex must then dissociate on dilution. Only polymers structurally similar to nucleic acids appear able to exploit this mechanism, highlighting their prevalence as informational macromolecules. The speed and volume of cycling could be critical determinants. Polymers will only accumulate if the rate of assembly (depending on cycling speed) exceeds the rate of fragmentation. The absolute number (rather than concentration) could be important. If the subsequent emergence of true lifeforms (defined as capable of self-replication and metabolism independent of physical cycling) requires one or more evolutionary jumps (e.g. auto-catalytic polymers, coding, metabolism and cellular inclusion) the likelihood of such rare events is in proportion to absolute number of prebiotic polymers. The emergence of life does not strictly require fast and large cycling, but such cycling might increase the likelihood.

Tidal cycling at the early terrestrial surface is favoured by a large and close Moon: the tidal periodicity in the Archaean is not known, but may have been fast. The presence of a large close satellite is not an absolute requirement. In the absence

of any moon the terrestrial tidal range (driven by Solar tides) is a significant fraction ($\sim 2/5$) of the lunar tide. Identical processes can operate with a tidal height range of 5 cm versus 5 m (although wave action at low tidal range would compromise the process).

The fast cycling requirement is further reduced if a mechanism exists to reassemble intact molecules from broken chains. Polymer scavenging (the recruitment of preformed fragments into a longer chain) affords such a mechanism. This possibility could favour fragment ligation as a primary target for nucleic acid (ribozyme-type) catalysis: both RNA and DNA ‘enzymes’ with ligase activity have been described (e.g. Cuenoud & Szostak (1995)).

To conclude, the picture painted here is of a steamy hot flat oceanscape, broken only by volcanic islands. Coastal erosion soon forms beaches, while a fast orbiting Moon, closer than today, drives regular tidal flooding of the shoreline. Rains and dews periodically inundate inland depressions. Evaporation then concentrates precursors in pools and sands, where partial dehydration favours aggregation and chemical linkage. Complex macromolecules are formed. Some are capable of template-directed assembly on concentration, but remain firmly locked together. Only molecules similar to nucleic acids dissociate on subsequent dilution, to allow another round to take place. Over time, with cyclic drying and dilution, these become more abundant.

Assembly was probably disordered at the beginning. The composition of new strands is dictated only by base-pairing efficacy and reactive substrate availability. Diverse structures accumulate, with many different sugars, nucleobases and non-classical linkages. Information content, however, can be conserved even if backbones and bases vary. Some sequences in these drying pools have unusually favourable properties – they are easily copied, relatively stable and perhaps they or their daughters have helpful enzymatic activity – these then accumulate at the expense of others (Darwinian selection), scavenging fragments of less-successful molecules.

If all of these processes operated, amplification of primitive nucleic acids was possibly not so much a limiting factor; indeed, the early seas could have teemed with these molecules. The next steps are perhaps more critical, including the evolution of coding, metabolism and a cellular environment. Tidal cycling could contribute to these evolutionary leaps. Long-chain hydrocarbons may have arisen in the Archaean; with wind, wave and tidal action, one imagines the formation of foams, froths and other micelles that accumulate on ocean margins. Further consideration of the turbulence of the prebiotic soup (itself influenced by tidal ebb and flow) may provide new pointers to prebiotic evolution.

Acknowledgements

The author is indebted to John O. Bishop for in-depth comments on the manuscript. Dieter Braun, Paul Knauth, Erik Kvale, Peter Olive, Ib Rasmussen, Matthias Tomczak and Jim Wessel Walker are thanked for further suggestions.

References

- Altman, S. (1990). Nobel lecture. Enzymatic cleavage of RNA by RNA. *Biosci. Rep.* **10**, 317–337.
- Arscott, P.G., Li, A.Z. & Bloomfield, V.A. (1990a). Condensation of DNA by trivalent cations. 1. Effects of DNA length and topology on the size and shape of condensed particles. *Biopolymers* **30**, 619–630.
- Arscott, P.G., Ma, C., Wenner, J. & Bloomfield, A.A. (1990b). DNA condensation by cobalt hexammine(III) in alcohol–water mixtures: dielectric constant and other solvent effects. *Biopolymers* **30**, 619–630.
- Auffinger, P., Bielecki, L. & Westhof, E. (2004). Anion binding to nucleic acids. *Structure* **12**, 379–388.
- Barringer, K.J., Orgel, L., Wahl, G. & Gingeras, T.R. (1990). Blunt-end and single-strand ligations by *Escherichia coli* ligase: influence on an *in vitro* amplification scheme. *Gene* **89**, 117–122.
- Bernal, J.D. (1960). Thermodynamics and kinetics of spontaneous generation. *Nature* **186**, 694–695.
- Bernal, J.D. (1949). The physical basis of life. *Proc. Phys. Soc. B.* **62**, 537–558.
- Bernal, J.D. (1961). The origin of life on the shores of the ocean: physical and chemical conditions determining first appearance of biological processes. In *Oceanography*, ed. Sears, M., pp. 95–118. American Association for the Advancement of Science, Washington, DC.
- Bernal, J.D. (1967). *The Origin of Life*. Weidenfeld and Nicolson, London.
- Bi, W. & Stambrook, P.J. (1997). CCR: a rapid and simple approach for mutation detection. *Nucleic Acids Res.* **25**, 2949–2951.
- Bloomfield, V.A. (1991). Condensation of DNA by multivalent cations: considerations on mechanism. *Biopolymers* **31**, 1481.
- Blum, H.F. (1957). On the origin of self-replicating systems. In *Rhythmic and Synthetic Processes in Growth*, ed. Rudnick, D., pp. 155–170. Princeton University Press, Princeton, NJ.
- Blum, H.F. (1962). On the origin and evolution of living machines. *Amer. Scientist* **47**, 474–501.
- Blume, S.W. *et al.* (1999). The integral divalent cation within the intermolecular purine*purine.pyrimidine structure: a variable determinant of the potential for and characteristics of the triple helical association. *Nucleic Acids Res.* **27**, 695–702.
- Bohler, C., Nielsen, P.E. & Orgel, L.E. (1995). Template switching between PNA and RNA oligonucleotides. *Nature* **376**, 578–581.
- Brack, A. (1987). Selective emergence and survival of early polypeptides in water. *Origins Life Evol. Biosphere* **17**, 367–379.
- Braun, D., Goddard, N.L. & Libchaber, A. (2003). Exponential DNA replication by laminar convection. *Phys. Rev. Lett.* **91**, 1–4.
- Braun, D. & Libchaber, A. (2002). Trapping of DNA by thermophoretic depletion and convection. *Phys. Rev. Lett.* **89**, 1–4.
- Braun, D. & Libchaber, A. (2004). Thermal force approach to molecular evolution. *Phys. Biol.* **1**, 1–8.
- Breaker, R.R. (1997). DNA enzymes. *Nat. Biotechnol.* **15**, 427–431.
- Brush, S.G. (1986). Early history of selenogony. In *Origin of the Moon*, eds Hartmann, W.K., Phillips, R.J. & Taylor, G.J., pp. 3–15. Lunar and Planetary Institute Houston, TX.
- Burak, Y., Ariel, G. & Andelman, D. (2003). Onset of DNA aggregation in the presence of monovalent and multivalent counterions. *Biophys. J.* **85**, 2100–2110.
- Butikov, E.I. (2002). A dynamical picture of the oceanic tides. *Amer. J. Phys.* **70**, 1001–1011.
- Cairns-Smith, A.G. (1982). *Genetic Takeover and the Mineral Origins of Life*. Cambridge University Press, Cambridge.
- Cairns-Smith, A.G. (2001). The origin of life: clays. In *Frontiers of Life*, vol. 1, eds Baltimore, D., Dulbecco, R., Jacob, F. & Levi-Montalcini, R., pp. 169–192. Academic Press, London.
- Calvin, M. (1969). *Chemical Evolution*. Oxford University Press, Oxford.
- Casey, J. & Davidson, N. (1976). Rates of formation and thermal stabilities of RNA–DNA and DNA–DNA duplexes at high concentrations of formamide. *Nucleic Acids Res.* **4**, 1539–1552.
- Cheng, A.J. & Van Dyke, M.W. (1993). Monovalent cation effects on intermolecular purine–purine–pyrimidine triple-helix formation. *Nucleic Acids Res.* **21**, 5630–5635.
- Clausen-Schaumann, H., Rief, M., Tolksdorf, C. & Gaub, H.E. (2000). Mechanical stability of single DNA molecules. *Biophys. J.* **78**, 1997–2007.
- Cleaves, H.J. (2002). The reactions of nitrogen heterocycles with acrolein: scope and prebiotic significance. *Astrobiology* **2**, 403–415.
- Corliss, J.B. *et al.* (1979). Submarine thermal springs on the Galápagos rift. *Science* **203**, 1073–1083.
- Cox, R.A. & Kanagalingam, K. (1968). A spectrophotometric study of the denaturation of deoxyribonucleic acid in the presence of urea or formaldehyde and its relevance to the secondary structure of single-stranded polynucleotides. *Biochem. J.* **108**, 599–610.
- Cuenoud, B. & Szostak, J.W. (1995). A DNA metalloenzyme with DNA ligase activity. *Nature* **375**, 611–614.
- Darwin, G.H. (1879). On the precession of a various spheroid, and on the remote history of the Earth. *Phil. Trans. R. Soc. Lond.* **170**, 447–538.
- Dickey, J.O. *et al.* (1994). Lunar laser ranging: a continuing legacy of the Apollo program. *Science* **265**, 482–490.
- Ding, P.Z., Kawamura, K. & Ferris, J.P. (1996). Oligomerization of uridine phosphorimidazolides on montmorillonite: a model for the prebiotic synthesis of RNA on minerals. *Origins Life Evol. Biosphere* **26**, 151–171.
- Dugaiczky, A., Boyer, H.W. & Goodman, H.M. (1975). Ligation of EcoRI endonuclease-generated DNA fragments into linear and circular structures. *J. Mol. Biol.* **96**, 171–184.
- Dworkin, J.P., Lazcano, A. & Miller, S.L. (2003). The roads to and from the RNA world. *J. Theor. Biol.* **222**, 127–134.
- Egholm, M., Buchardt, O., Christensen, L., Behrens, C., Freier, S.M., Driver, D.A. *et al.* (1993). PNA hybridizes to complementary oligonucleotides obeying the Watson–Crick hydrogen-bonding rules. *Nature* **365**, 566–568.
- Eickbush, T.H. & Moudrianakis, E.N. (1978). The compaction of DNA helices into either continuous supercoils or folded-fiber rods and toroids. *Cell* **13**, 295–306.
- Eli, P., Ariyama, T., Nishigaki, K. & Husimi, Y. (1999). Characterization of imidazole as a DNA denaturant by using TGGE of PCR products from a random pool of DNA. *J. Biochem. (Tokyo)* **125**, 790–794.
- Ertem, G. (2004). Montmorillonite, oligonucleotides, RNA and origin of life. *Origins Life Evol. Biosphere* **34**, 549–570.
- Ertem, G. & Ferris, J.P. (1996). Synthesis of RNA oligomers on heterogeneous templates. *Nature* **379**, 238–240.
- Ertem, G. & Ferris, J.P. (2000). Sequence- and regio-selectivity in the montmorillonite-catalyzed synthesis of RNA. *Origins Life Evol. Biosphere* **30**, 411–422.
- Ferris, J.P. (1987). Prebiotic synthesis – problems and challenges. *Cold Spring Harbour Symp. Quant. Biol.* **52**, 29–35.
- Ferris, J.P. (1992). Chemical markers of prebiotic chemistry in hydrothermal systems. *Origins Life Evol. Biosphere* **22**, 109–242.
- Ferris, J.P. & Hagan, W.J. Jr. (1984). HCN and chemical evolution: the possible role of cyano compounds in prebiotic synthesis. *Tetrahedron* **40**, 1093–1120.
- Ferris, J.P., Hill, A. Jr., Liu, R. & Orgel, L.E. (1996). Synthesis of long prebiotic oligomers on mineral surfaces. *Nature* **381**, 59–61.
- Ferris, J.P., Yanagawa, H. & Hagan, W.J. Jr. (1983). Prebiotic synthesis and reactions of nucleosides and nucleotides. *Adv. Space Sci.* **3**, 61–68.
- Floris, R., Scaggiante, B., Manzini, G., Quadrioglio, F. & Xodo, L.E. (1999). Effect of cations on purine.purine.pyrimidine triple helix formation in mixed-valence salt solutions. *Eur. J. Biochem.* **260**, 801–809.
- Fox, S.W. & Harada, K. (1958). Thermal copolymerization of amino acids to a product resembling protein. *Science* **128**, 1214.
- Franklin, R. & Gosling, R. (1953). The structure of sodium thymonucleate fibres: II. The influence of water content. *Acta Crystal.* **6**, 673–677.
- Fuller, W.D., Sanchez, R.A. & Orgel, L.E. (1972). Studies in prebiotic synthesis. VI. Synthesis of purine nucleosides. *J. Mol. Biol.* **67**, 25–33.
- Gao, K. & Orgel, L.E. (2000). Polyphosphorylation and non-enzymatic template-directed ligation of oligonucleotides. *Origins Life Evol. Biosphere* **30**, 45–51.
- Golos, B., Dzik, J.M., Kazimierczuk, Z., Ciesla, J., Zielinski, Z., Jankowska, J. *et al.* (2001). Interaction of thymidylate synthase with

- the 5'-thiophosphates, 5'-dithiophosphates, 5'-H-phosphonates and 5'-S-thiosulfates of 2'-deoxyuridine, thymidine and 5-fluoro-2'-deoxyuridine. *Biol. Chem.* **382**, 1439–1445.
- Gosule, L.C. & Schellman, J.A. (1976). Compact form of DNA induced by spermidine. *Nature* **259**, 333–335.
- Gulick, A. (1955). Phosphorus as a factor in the origin of life. *Amer. Sci.* **43**, 479–489.
- Haldane, J.B.S. (1929). Science and human life. *Rational. Ann.* **148**, 3.
- Hamaguchi, K. & Geiduschek, E.P. (1962). The effect of electrolytes on the stability of the deoxyribonucleate helix. *J. Amer. Chem. Soc.* **84**, 1329–1338.
- Hames, B.D. & Higgins, S.J. (1985). *Nucleic Acid Hybridisation: A Practical Approach*. IRL Press, Oxford.
- Hartmann, W.K. (1980). Dropping stones in magma oceans – effects of early lunar cratering. In *Proc. Conf. Lunar Highlands Crust*, eds Papike, J.J. & Merrill, R.B., pp. 155–171. Pergamon Press, New York.
- Hartmann, W.K. (1999). *Moons and Planets*. Wadsworth, Belmont, CA.
- Haug, P.C. (1968). Effect of guanidinium ion on DNA denaturation and renaturation. *Biochem. Biophys. Res. Commun.* **33**, 384–390.
- Hull, D.E. (1960). Thermodynamics and kinetics of spontaneous generation. *Nature* **186**, 693–694.
- Huxley, T.H. (1870). Biogenesis and abiogenesis. *Address to the British Association for the Advancement of Science*. Liverpool.
- Imai, E., Honda, H., Hatori, K., Brack, A. & Matsuno, K. (1999). Elongation of oligopeptides in a simulated submarine hydrothermal system. *Science* **283**, 831–833.
- AQ3 Jacobson, H. & Stockmayer, W.H. (1950). Intramolecular reaction in polycondensations. I. The theory of linear systems. *J. Chem. Phys.* **18**, 1600–1606.
- Jaroszewski, J.W., Clausen, V., Cohen, J.S. & Dahl, O. (1996). NMR investigations of duplex stability of phosphorothioate and phosphorodithioate DNA analogues modified in both strands. *Nucleic Acids Res.* **24**, 829–834.
- Joyce, G.F., Schwartz, A.W., Miller, S.L. & Orgel, L.E. (1987). The case for an ancestral genetic system involving simple analogues of the nucleotides. *Proc. Natl. Acad. Sci. USA* **84**, 4398–4402.
- Kasting, J.F. & Ackerman, T.P. (1986). Climatic consequences of very high carbon dioxide levels in the Earth's early atmosphere. *Science* **234**, 1383–1385.
- Keefe, A.D. & Miller, S.L. (1995). Are polyphosphates or phosphate esters prebiotic reagents? *J. Mol. Evol.* **41**, 693–702.
- Knauth, L.P. (1998). Salinity history of the Earth's early ocean. *Nature* **395**, 555.
- Knauth, L.P. (2005). Temperature and salinity history of the Precambrian ocean: implications for the course of microbial evolution. *Palaeogeog. Palaeoclimatol. Palaeoecol.* **219**, 53–69.
- AQ4 Kourilsky, P., Leidner, J. & Tremblay, G.Y. (1971). DNA–DNA hybridization on filters at low temperature in the presence of formamide or urea. *Biochimie* **53**, 1111–1114.
- Kourilsky, P., Manteuil, S., Zamansky, M.H. & Gros, F. (1970). DNA–RNA hybridization at low temperature in the presence of urea. *Biochem. Biophys. Res. Commun.* **41**, 1080–1087.
- Krasinsky, G.A. (2002). Dynamical history of the Earth–Moon system. *Celest. Mech. Dynam. Astron.* **84**, 27–55.
- Krishnan, M., Ugaz, V.M. & Burns, M.A. (2002). PCR in a Rayleigh–Benard convection cell. *Science* **298**, 793.
- Kvale, E.P., Johnson, H.W., Sonett, C.P., Archer, A.W. & Zawistoski, A. (1999). Calculating Lunar retreat rates using tidal rhythmites. *J. Sedimentary Res.* **69**, 1154–1168.
- Lambeck, K. (1978). The Earth's paleorotation. In *Tidal Friction and the Earth's Rotation*, eds Brosche, P. & Sundermann, J., pp. 145–153. Springer, Berlin.
- Lambeck, K. (1980). *The Earth's Variable Rotation: Geophysical Causes and Consequences*. Cambridge University Press, Cambridge.
- Larralde, R., Robertson, M.P. & Miller, S.L. (1995). Rates of decomposition of ribose and other sugars: implications for chemical evolution. *Proc. Natl. Acad. Sci. USA* **92**, 8158–8160.
- Lathe, R. (1985). Synthetic oligonucleotide probes deduced from amino acid sequence data. Theoretical and practical considerations. *J. Mol. Biol.* **183**, 1–12.
- Lathe, R. (2004). Fast tidal cycling and the origin of life. *Icarus* **168**, 18–22.
- Lathe, R. (2005). Early tides. *Icarus*, submitted.
- AQ4 Legerski, R.J. & Robberson, D.L. (1985). Analysis and optimization of recombinant DNA joining reactions. *J. Mol. Biol.* **181**, 297–312.
- Levy, M. & Miller, S.L. (1998). The stability of the RNA bases: implications for the origin of life. *Proc. Natl. Acad. Sci. USA* **95**, 7933–7938.
- Liu, R. & Orgel, L.E. (1998). Polymerization on the rocks: beta-amino acids and arginine. *Origins Life Evol. Biosphere* **28**, 245–257.
- Lohrmann, R., Bridson, P.K. & Orgel, L.E. (1980). Efficient metal-ion catalyzed template-directed oligonucleotide synthesis. *Science* **208**, 1464–1465.
- Lohrmann, R. & Orgel, L.E. (1971). Urea-inorganic phosphate mixtures as prebiotic phosphorylating agents. *Science* **171**, 490–494.
- Luther, A., Brandsch, R. & von Kiedrowski, G. (1998). Surface-promoted replication and exponential amplification of DNA analogues. *Nature* **396**, 245–248.
- MacLeod, G., McKeown, C., Hall, A.J. & Russell, M.J. (1994). Hydrothermal and oceanic pH conditions of possible relevance to the origin of life. *Origins Life Evol. Biosphere* **24**, 19–41.
- Maher, K.A. & Stevenson, D.J. (1988). Impact frustration of the origin of life. *Nature* **331**, 612–614.
- McConaughy, B.L., Laird, C.D. & McCarthy, B.J. (1969). Nucleic acid reassociation in formamide. *Biochemistry* **8**, 3289–3295.
- Miller, S.L. (1953). A production of amino acids under possible primitive Earth conditions. *Science* **117**, 528–529.
- Miller, S.L. & Bada, J.L. (1988). Submarine hot springs and the origin of life. *Nature* **334**, 609–611.
- Miller, S.L. & Lazcano, A. (1995). The origin of life – did it occur at high temperatures? *J. Mol. Evol.* **41**, 689–692.
- Miller, S.L. & Urey, H.C. (1959). Organic compound synthesis on the primitive Earth. *Science* **130**, 245–251.
- Mizutani, T. & Narihara, T. (1982). Adsorption chromatography of nucleic acids on siliconized porous glass. *Nucleic Acids Symp. Ser.* **1982**, 127–130.
- AQ5 Mullis, K., Faloona, F., Scharf, S., Saiki, R., Horn, G. & Erlich, H. (1986). Specific enzymatic amplification of DNA in vitro: the polymerase chain reaction. *Cold Spring Harbour Symp. Quant. Biol.* **51(1)**, 263–273.
- Musso, M., Thomas, T., Shirahata, A., Sigal, L.H., Van Dyke, M.W. & Thomas, T.J. (1997). Effects of chain length modification and bis(ethyl) substitution of spermine analogs on purine–purine–pyrimidine triplex DNA stabilization, aggregation, and conformational transitions. *Biochemistry* **36**, 1441–1449.
- Musso, M. & Van Dyke, M.W. (1995). Polyamine effects on purine–purine–pyrimidine triple helix formation by phosphodiester and phosphorothioate oligodeoxyribonucleotides. *Nucleic Acids Res.* **23**, 2320–2327.
- Nakano, S., Fujimoto, M., Hara, H. & Sugimoto, N. (1999). Nucleic acid duplex stability: influence of base composition on cation effects. *Nucleic Acids Res.* **27**, 2957–2965.
- Naylor, R. & Gilham, P.T. (1966). Studies on some interactions and reactions of oligonucleotides in aqueous solution. *Biochemistry* **5**, 2722–2728.
- Nelson, K.E., Levy, M. & Miller, S.L. (2000). Peptide nucleic acids rather than RNA may have been the first genetic molecule. *Proc. Natl. Acad. Sci. USA* **97**, 3868–3871.
- Nelson, K.E., Robertson, M.P., Levy, M. & Miller, S.L. (2001). Concentration by evaporation and the prebiotic synthesis of cytosine. *Origins Life Evol. Biosphere* **31**, 221–229.
- Nielsen, P.E., Egholm, M., Berg, R.H. & Buchardt, O. (1991). Sequence-selective recognition of DNA by strand displacement with a thymine-substituted polyamide. *Science* **254**, 1497–1500.
- Nisbet, E.G. & Sleep, N.H. (2001). The habitat and nature of early life. *Nature* **409**, 1083–1091.
- Ogata, N. & Miura, T. (1997). Genetic information 'created' by archaeobacterial DNA polymerase. *Biochem. J.* **324(2)**, 667–671.

- Oparin, A.I. (1953). *The Origin of Life*, 2nd edn. Academic Press, New York.
- Orgel, L.E. (1998). Polymerization on the rocks: theoretical introduction. *Origins Life Evol. Biosphere* **28**, 227–234.
- Oro, J. (1961). Mechanism of synthesis of adenine from hydrogen cyanide under possible primitive Earth conditions. *Nature* **191**, 1193–1194.
- Ott, G.S., Ziegler, R. & Bauer, W.R. (1975). The DNA melting transition in aqueous magnesium salt solutions. *Biochemistry* **14**, 3431–3438.
- Panella, G. (1975). Paleontological clocks and the history of the Earth's rotation. In *Growth Rhythms and the History of the Earth's Rotation*, eds Rosenberg, G.D. & Runcorn, S.K., pp. 253–284. Wiley, Chichester.
- Parnell, J. (2004). Mineral radioactivity in sands as a mechanism for fixation of organic carbon on the early Earth. *Origins Life Evol. Biosphere* **34**, 553–547.
- Peracchi, A. (1999). Origins of the temperature dependence of hammerhead ribozyme catalysis. *Nucleic Acids Res.* **27**, 2875–2882.
- Piskur, J. & Rupprecht, A. (1995). Aggregated DNA in ethanol solution. *FEBS Lett.* **375**, 174–178.
- Plum, G.E., Pilch, D.S., Singleton, S.F. & Breslauer, K.J. (1995). Nucleic acid hybridization: triplex stability and energetics. *Ann. Rev. Biophys. Biomol. Struct.* **24**, 319–350.
- Reysenbach, A.L. & Shock, E. (2002). Merging genomes with geochemistry in hydrothermal ecosystems. *Science* **296**, 1077–1082.
- Rose, K., Mason, J.O. & Lathe, R. (2002). Hybridization parameters revisited: solutions containing SDS. *Biotechniques* **33**, 54–58.
- Saminathan, M., Antony, T., Shirahata, A., Sigal, L.H., Thomas, T. & Thomas, T.J. (1999). Ionic and structural specificity effects of natural and synthetic polyamines on the aggregation and resolubilization of single-, double-, and triple-stranded DNA. *Biochemistry* **38**, 3821–3830.
- Sanchez, R.A., Ferris, J.P. & Orgel, L.E. (1967). Studies in prebiotic synthesis. II. Synthesis of purine precursors and amino acids from aqueous hydrogen cyanide. *J. Mol. Biol.* **30**, 223–253.
- Sanchez, R.A. & Orgel, L.E. (1970). Studies in prebiotic synthesis. V. Synthesis and photoanomerization of pyrimidine nucleosides. *J. Mol. Biol.* **47**, 531–543.
- Sandars, P.G.H. (2005). Chirality in the RNA world and beyond. *Int. J. Astrobiol.*, this volume.
- Schildkraut, C. & Lifson, S. (1965). Dependence of the melting temperature of DNA on the salt concentration. *Biopolymers* **3**, 195–208.
- Schlesinger, G. & Miller, S.L. (1983). Prebiotic synthesis in atmospheres containing CH₄, CO, and CO₂. II. Hydrogen cyanide, formaldehyde and ammonia. *J. Mol. Evol.* **19**, 383–390.
- Schneider, B., Patel, K. & Berman, H.M. (1998). Hydration of the phosphate group in double-helical DNA. *Biophys. J.* **75**, 2422–2434.
- Schoning, K., Scholz, P., Guntha, S., Wu, X., Krishnamurthy, R. & Eschenmoser, A. (2000). Chemical etiology of nucleic acid structure: the alpha-threofuranosyl-(3'→2') oligonucleotide system. *Science* **290**, 1347–1351.
- Schramm, G., Grotzsch, W. & Pollman, W. (1962). Nonenzymatic synthesis of polysaccharides, nucleosides, nucleic acids and the origin of self-reproducing systems. *Angew. Chem. (Int. English. Edn)* **1**, 1–7.
- Scrutton, C.T. (1978). Periodic growth features in fossil organisms and the length of the day and month. In *Tidal Friction and the Earth's Rotation*, eds Brosche, P. & Sundermann, J., pp. 154–196. Springer, Berlin.
- Shapiro, R. (1988). Prebiotic ribose synthesis: a critical analysis. *Origins Life Evol. Biosphere* **18**, 71–85.
- Shapiro, R. (1995). The prebiotic role of adenine: a critical analysis. *Orig. Life Evol. Biosph.* **25**, 83–98.
- Shapiro, R. (1999). Prebiotic cytosine synthesis: a critical analysis and implications for the origin of life. *Proc. Natl. Acad. Sci. USA* **96**, 4396–4401.
- Shapiro, R. (2002). Comments on 'concentration by evaporation and the prebiotic synthesis of cytosine'. *Origins Life Evol. Biosphere* **32**, 275–278.
- Shock, E.L. (1996). Hydrothermal systems as environments for the emergence of life. *Ciba Found. Symp.* **202**, 40–52.
- Singleton, S.F. & Dervan, P.B. (1993). Equilibrium association constants for oligonucleotide-directed triple helix formation at single DNA sites: linkage to cation valence and concentration. *Biochemistry* **32**, 13 171–13 179.
- Sonett, C.P., Kvale, E.P., Zakharian, A., Chan, M.A. & Dernko, T.M. (1996). Late Proterozoic and Paleozoic tides, retreat of the Moon, and rotation of the Earth. *Science* **273**, 100–104.
- Takagi, Y. & Taira, K. (2002). Analyses of kinetic solvent isotope effects of a hammerhead ribozyme reaction in NH₄⁺ and Li⁺ ions. *Nucleic Acids Res. Suppl.* 273–274.
- Thomas, C.A.J. & Dancis, B.M. (1973). Ring stability. *J. Mol. Biol.* **77**, 43–55.
- Vanyo, J.P. & Awramik, S.M. (1982). Length of day and obliquity of the ecliptic 850 MA ago: preliminary results of a stromatolite growth model. *Geophys. Rev. Lett.* **9**, 1125–1128.
- Varga, P., Rybicki, K.R. & Denis, C. (2005). Comment on the paper 'Fast tidal cycling and the origin of life'. *Icarus*, submitted.
- Vogelstein, B. & Gillespie, D. (1979). Preparative and analytical purification of DNA from agarose. *Proc. Natl. Acad. Sci. USA* **76**, 615–619.
- von Kiedrowski, G., Wlotzka, B. & Helbing, J. (1989). Sequence dependence of template-directed syntheses of hexadeoxynucleotide derivatives with 3'-5' pyrophosphate linkages. *Angew. Chem. (Int. English. Edn)* **28**, 1235–1237.
- Wachtershauser, G. (1988). An all-purine precursor of nucleic acids. *Proc. Natl. Acad. Sci. USA* **85**, 1134–1135.
- Walker, J.C. & Zahnle, K.J. (1986). Lunar nodal tide and distance to the Moon during the Precambrian. *Nature* **320**, 600–602.
- Weber, A.L., Caroon, J.M., Warden, J.T., Lemmon, R.M. & Calvin, M. (1977). Simultaneous peptide and oligonucleotide formation in mixtures of amino acid, nucleoside triphosphate, imidazole, and magnesium ion. *Biosystems* **8**, 277–286.
- Westhof, E. (1988). Water: an integral part of nucleic acid structure. *Ann. Rev. Biophys. Biophys. Chem.* **17**, 125–144.
- Wetmur, J.G. (1991). DNA probes: applications of the principles of nucleic acid hybridization. *Crit. Rev. Biochem. Mol. Biol.* **26**, 227–259.
- Wetmur, J.G. & Davidson, N. (1986). Kinetics of renaturation of DNA. *J. Mol. Biol.* **31**, 349–370.
- Williams, A.P., Longfellow, C.E., Freier, S.M., Kierzek, R. & Turner, D.H. (1989). Laser temperature-jump, spectroscopic, and thermodynamic study of salt effects on duplex formation by dGCATGC. *Biochemistry* **28**, 4283–4291.
- Williams, G.E. (1997). Precambrian length of day and the validity of tidal rhythmite paleotidal values. *Geophys. Rev. Lett.* **21**, 421–424.
- Williams, G.E. (2000). Geological constraints on the precambrian history of Earth's rotation and the Moon's orbit. *Rev. Geophys.* **38**, 37–59.
- Woese, C.R. (2000). Interpreting the universal phylogenetic tree. *Proc. Natl. Acad. Sci. USA* **97**, 8392–8396.
- Woese, C.R., Kandler, O. & Wheelis, M.L. (1990). Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proc. Natl. Acad. Sci. USA* **87**, 4576–4579.
- Wu, D.Y. & Wallace, R.B. (1989). The ligation amplification reaction (LAR)–amplification of specific DNA sequences using sequential rounds of template-dependent ligation. *Genomics* **4**, 560–569.
- Xu, Y. & Kool, E.T. (1999). High sequence fidelity in a non-enzymatic DNA autoligation reaction. *Nucleic Acids Res.* **27**, 875–881.
- Yamagata, Y., Watanabe, H., Saitoh, M. & Namba, T. (1991). Volcanic production of polyphosphates and its relevance to prebiotic evolution. *Nature* **352**, 516–519.

AQ4

AQ4